

**ELECTROMAGNETIC SPINOR AND WAVE  
FUNCTIONS IN MINKOWSKI SPACETIME**

**A. Lucas-Bravo**

UPIITA, Instituto Politécnico Nacional, Av. IPN 2580  
Col. Barrio la Laguna 07340, CDMX, México  
andreslbravo@yahoo.com.mx

**J. López-Bonilla, R. López-Vázquez**

Instituto Politécnico Nacional  
Col. Lindavista CP 07738, CDMX, México  
jlopezb@ipn.mx, gravitonramiro@hotmail.com

Received December 7, 2022

**Abstract**

We show how generate the electromagnetic spinor verifying the Maxwell equations in vacuum; we also exhibit the Whittaker and Bateman wave functions and their connection with electromagnetic fields in Minkowski geometry.

**Keywords:** Maxwell spinor, Riemann-Silberstein vector, Maxwell equations, Wave functions, Quaternions.

All electromagnetic information is contained in the Faraday's skew-symmetric tensor [1-3]:

$$(F^{\mu\nu}) = \begin{pmatrix} 0 & -E_1 & -E_2 & -E_3 \\ E_1 & 0 & -cB_3 & cB_2 \\ E_2 & cB_3 & 0 & -cB_1 \\ E_3 & -cB_2 & cB_1 & 0 \end{pmatrix}, \quad (1)$$

where  $\vec{E} = (E_1, E_2, E_3)$  and  $\vec{B} = (B_1, B_2, B_3)$  are the electric and magnetic fields expressed in the MKS system of units, respectively, verifying the Maxwell equations in empty spacetime:

$$\vec{\nabla} \cdot \vec{E} = \vec{\nabla} \cdot \vec{B} = 0, \quad \vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}, \quad \vec{\nabla} \times \vec{B} = \frac{1}{c^2} \frac{\partial \vec{E}}{\partial t}. \quad (2)$$

If we introduce the Riemann [4]-Silberstein [5, 6] complex vector (thus named by Bialynicki-Birula [7]) [8]:

$$\vec{F} = c \vec{B} + i \vec{E}, \quad (3)$$

then (2) are equivalent to two complex equations [9]:

$$\vec{\nabla} \cdot \vec{F} = 0, \quad \vec{\nabla} \times \vec{F} + \frac{i}{c} \frac{\partial}{\partial t} \vec{F} = \vec{0}, \quad (4)$$

and the corresponding spinorial form is given by [10-14]:

$$\partial_{B\dot{C}} \varphi^{AB} = 0, \quad (5)$$

involving the symmetric Maxwell spinor. It is clear that, without boundary conditions, the equations (5) have many solutions, thus it is important to have a method to construct them.

On the other hand, we know that a unitary quaternion [15-17] generates rotations in three and four dimensions [18-20], but, what happens with a null quaternion? Here we consider the following case:

$$\mathbf{A} = -\sin \tau - i \mathbf{I} + \cos \tau \mathbf{K} \quad \therefore \quad \mathbf{A}\bar{\mathbf{A}} = (\sin \tau)^2 + i^2 + (\cos \tau)^2 = 0, \quad (6)$$

with its associated 2x2 complex matrix:

$$\tilde{M} = \begin{pmatrix} ie^{i\tau} & 1 \\ 1 & -ie^{-i\tau} \end{pmatrix}. \quad (7)$$

Now the interesting result is that (7) allows construct solutions for (5), in fact:

$$\begin{aligned} (\varphi^{AB}) &= \frac{1}{2} \int_0^{2\pi} \tilde{M} G(u, v, \tau) d\tau, \quad u = x \cos \tau + y \sin \tau + i z, \\ v &= x \sin \tau - y \cos \tau + c t, \end{aligned} \quad (8)$$

where  $G$  is an arbitrary function of its arguments; hence (8) is a factory to elaborate solutions for the Maxwell equations in vacuum, that is:

$$\begin{aligned} \varphi^{11} &= \frac{i}{2} \int_0^{2\pi} e^{i\tau} G d\tau, \quad \varphi^{12} = \varphi^{21} = \frac{1}{2} \int_0^{2\pi} G d\tau, \\ \varphi^{22} &= -\frac{i}{2} \int_0^{2\pi} e^{-i\tau} G d\tau. \end{aligned} \quad (9)$$

We have the expressions for the spinor covariant derivative [12]:

$$\begin{aligned} \partial_{1i} &= \frac{1}{\sqrt{2}} \left( \frac{\partial}{c \partial t} + \frac{\partial}{\partial z} \right), \quad \partial_{2i} = \overline{\partial_{1\bar{2}}} = \frac{1}{\sqrt{2}} \left( \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right), \\ \partial_{2\bar{2}} &= \frac{1}{\sqrt{2}} \left( \frac{\partial}{c \partial t} - \frac{\partial}{\partial z} \right), \end{aligned} \quad (10)$$

then it is simple to prove the relations:

$$\begin{aligned} \partial_{1i} G &= \frac{1}{\sqrt{2}} \left( \frac{\partial G}{\partial v} + i \frac{\partial G}{\partial u} \right) = i e^{-i\tau} \partial_{2i} G, \quad \partial_{2\bar{2}} G = \frac{1}{\sqrt{2}} \left( \frac{\partial G}{\partial v} - i \frac{\partial G}{\partial u} \right) = \\ &= -i e^{i\tau} \partial_{1\bar{2}} G, \end{aligned} \quad (11)$$

thus with (9) and (11) is immediate verify (5), *q.e.d.*

The components of the Riemann-Silberstein complex vector (3) are given by:

$$F_1 = i(\varphi^{22} - \varphi^{11}) = \int_0^{2\pi} G \cos \tau d\tau, \quad F_2 = -(\varphi^{22} + \varphi^{11}) = \int_0^{2\pi} G \sin \tau d\tau, \quad F_3 = 2i\varphi^{12} = i \int_0^{2\pi} G d\tau,$$

in agreement with the result of Bateman [9]:

$$\vec{F} = \int_0^{2\pi} G(x \cos \tau + y \sin \tau + i z, x \sin \tau - y \cos \tau + c t, \tau) \vec{R} d\tau, \\ \vec{R} = (\cos \tau, \sin \tau, i). \quad (12)$$

We have the valuable theorem of I. Robinson and J. L. Synge [2, 22-25]:

“Every solution of the vacuum Maxwell equations in Minkowski spacetime can be written in terms of two real wave functions and a constant real bivector”, (13)

and the corresponding Faraday tensor is given by the expression:

$$F_{\mu\nu} = (H_\mu{}^\lambda U_{,\lambda} + {}^*H_\mu{}^\lambda V_{,\lambda})_{,\nu} - (H_\nu{}^\lambda U_{,\lambda} + {}^*H_\nu{}^\lambda V_{,\lambda})_{,\mu}, \quad (14)$$

where  $\square U = \square V = 0$  and  ${}^*H_{\mu\nu}$  is the dual of the constant skew-symmetric tensor  $H_{\mu\nu}$ .

The first person to suggest that a vacuum electromagnetic field could be constructed from only two real wave functions was Whittaker [23, 24, 26-29]: He showed explicitly that the Liénard-Wiechert field of an accelerating point charge could be derived from a pair of wave functions, and he calculated these functions. With the constant tensor:

$$(H^{\mu\nu}) = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \quad \therefore \quad ({}^*H^{\mu\nu}) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (15)$$

the relations (1) and (14) imply the following expressions of Whittaker [26]:

$$E_1 = \frac{\partial^2 U}{\partial x \partial z} + \frac{1}{c} \frac{\partial^2 V}{\partial y \partial t}, \quad E_2 = \frac{\partial^2 U}{\partial y \partial z} - \frac{1}{c} \frac{\partial^2 V}{\partial x \partial t}, \quad E_3 = \frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2}, \quad (16)$$

$$c B_1 = -\frac{\partial^2 V}{\partial x \partial z} + \frac{1}{c} \frac{\partial^2 U}{\partial y \partial t}, \quad c B_2 = -\frac{\partial^2 V}{\partial y \partial z} - \frac{1}{c} \frac{\partial^2 U}{\partial x \partial t}, \quad c B_3 = \frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2}.$$

Now we need a systematic method to construct the scalar wave functions  $U$  and  $V$ , in fact [26]:

$$U = \int_0^\pi \int_0^{2\pi} f(x \sin u \cos v + y \sin u \sin v + z \cos u + c t, u, v) du dv, \quad (17)$$

$$V = \int_0^\pi \int_0^{2\pi} g(x \sin u \cos v + y \sin u \sin v + z \cos u + c t, u, v) du dv,$$

where  $f$  and  $g$  are arbitrary functions of their arguments. Bateman [9, 30] showed an alternative approach to find wave functions in four dimensions:

$$W = \int_0^{2\pi} G(x \cos \tau + y \sin \tau + i z, x \sin \tau - y \cos \tau + c t, \tau) d\tau, \quad (18)$$

being  $G$  an arbitrary function. Finally, we comment that Whittaker [9, 31] proved that the function:

$$w = \int_0^{2\pi} h(z + i x \cos \alpha + i y \sin \alpha, \alpha) d\alpha, \quad (19)$$

for  $h$  an arbitrary function, satisfies the Laplace equation  $\nabla^2 w = 0$ .

## References

- [1] H. Minkowski, *Die grundgleichungen für die elektromagnetischen vorgänge in bewegten körpern*, Nach. Kgl. Gesell. Wiss. Göttingen (1908) 53-111.

- [2] J. L. Synge, *Relativity: the special theory*, North-Holland, Amsterdam (1965).
- [3] A. Hernández-Galeana, J. López-Bonilla, R. López-Vázquez, G. R. Pérez-Teruel, *Faraday tensor and Maxwell spinor*, Prespacetime Journal **6**, No. 2 (2015) 88-107.
- [4] B. Riemann, *Die partiellen Differential-Gleichungen der mathematische Physik*, Lecture notes edited by H. Weber **2**, Vieweg, Braunschweig (1901).
- [5] L. Silberstein, *Elektromagnetische Grundgleichungen in bivectorieller behandlung*, Ann. der Physik, Leipzig **22** (1907) 579-586.
- [6] L. Silberstein, *Nachtrag zur abhandlung über electromagnetische grundgleichungen in bivectorieller behandlung*, Ann. der Physik, Leipzig, **24** (1907) 783-784.
- [7] I. Bialynicki-Birula, *Photon wave function*, Prog. in Optics **36**, E. Wolf, Elsevier, Amsterdam (1996).
- [8] N. Hamdan, I. Guerrero-Moreno, J. López-Bonilla, L. Rosales-Roldán, *On the Faraday's complex vector*, The Icfai Univ. J. Phys. **1**, No. 3 (2008) 52-56.
- [9] H. Bateman, *Partial differential equations of mathematical physics*, Dover, New York (1944).
- [10] W. Rindler, *What are spinors?*, Am. J. Phys. **34**, No. 10 (1966) 937-942.
- [11] J. Hladik, *Spinors in Physics*, Springer, New York (1999).
- [12] P. O'Donnell, *Introduction to 2-spinors in general relativity*, World Scientific, Singapore (2003).
- [13] L. Ioan-Piscoran, J. López-Bonilla, R. López-Vázquez, *Spinorial and quaternionic forms of Maxwell equations in vacuo*, The SciTech, J. Sci. & Tech. **2**, No. 2 (2013) 29-32.
- [14] D. S. Kulyabov, A. V. Korolkova, L. A. Sevastianov, *Spinor representation of Maxwell equations*, J. of Physics: Conf. Series **788** (2017) 012025.
- [15] E. T. Whittaker, *The sequence of ideas in the discovery of quaternions*, Proc. Roy. Irish Acad. **A50**, No. 6 (1945) 93-98.
- [16] B. L. van der Waerden, *Hamilton's discovery of quaternions*, Maths. Magazine **49**, No. 5 (1976) 227-234.

- [17] T. Bannon, *The origin of quaternions*, Coll. Math. J. **46**, No. 1 (2015) 43-50.
- [18] J. López-Bonilla, J. Morales, G. Ovando, *Quaternions, 3-rotations and Lorentz transformations*, Indian J. Theor. Phys. **52**, No. 2 (2004) 91-96.
- [19] I. Guerrero-Moreno, J. López-Bonilla, L. Rosales, *Rotations in three and four dimensions via 2x2 complex matrices and quaternions*, The Icfai Univ. J. Phys. **1**, No. 2 (2008) 7-13.
- [20] P. A. M. Dirac, *Applications of quaternions to Lorentz transformations*, Proc. Roy. Irish Acad. **A50** (1945) 261-270.
- [21] J. L. Synge, *Relativity: the special theory*, North-Holland, Amsterdam (1965).
- [22] R. Penrose, *Zero-rest-mass fields including gravitation: Asymptotic behaviour*, Proc. Roy. Soc. London **A284** (1965) 159-203.
- [23] P. A. Hogan, *Newman-Penrose constants for zero-rest-mass fields in Minkowskian space-time*, J. Math. Phys. **28**, No. 9 (1987) 2089-2093.
- [24] P. A. Hogan, G. F. R. Ellis, *Aspects of asymptotic electromagnetic fields*, Ann. of Phys. **210** (1991) 178-237.
- [25] J. López-Bonilla, J. Morales, G. Ovando, J. J. Peña, *Lanczos potential for the Minkowski space*, Indian J. Phys. **B74**, No. 5 (2000) 393-395.
- [26] E. T. Whittaker, *On an expression of the electromagnetic field due to electrons by means of two scalar potential functions*, Proc. London Math. Soc. Series II, **1** (1904) 367-372.
- [27] H. S. Ruse, *On Whittaker's electromagnetic scalar potentials*, Quart. J. Maths. **8**, No. 1 (1937) 148-160.
- [28] P. Hillion, *Hertz and Whittaker scalar potentials*, Fundamental J. Math. Phys. **1**, No. 2 (2011) 53-55.
- [29] Y. Friedman, *The wave-function description of the electromagnetic field*, J. of Phys: Conf. Series **437** (2013) 012018.
- [30] H. Bateman, *The solution of partial differential equations by means of definite integrals*, Proc. London Math. Soc. **1**, No. 2 (1904) 451-458.
- [31] E. T. Whittaker, *On the partial differential equations of mathematical physics*, Math. Ann. **57** (1903) 333-355.